



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1992, 3rd Annual JWP Conference

Apr 25th, 10:30 AM - 4:30 PM

The Effects of Acute Administration of the 5-HT Neurotoxin PCA on the Expression of Male Rat Copulatory Behavior: A Comparison with MDMA

William M. Struthers

Illinois Wesleyan University

Wayne Dornan, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

William M. Struthers and Wayne Dornan, Faculty Advisor, "The Effects of Acute Administration of the 5-HT Neurotoxin PCA on the Expression of Male Rat Copulatory Behavior: A Comparison with MDMA" (April 25, 1992). *John Wesley Powell Student Research Conference*. Paper 11.
<http://digitalcommons.iwu.edu/jwprc/1992/posters/11>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

**THE EFFECTS OF ACUTE ADMINISTRATION OF THE 5-HT NEUROTOXIN
PCA ON THE EXPRESSION OF MALE RAT COPULATORY BEHAVIOR:
A COMPARISON WITH MDMA**

William M. Struthers, Dept. of Psychology, IWU
Dr. Wayne Dornan*

3,4 Metylenedioxymethamphetamine (MDMA) is a potent neurotoxin which produces 5-HT nerve terminal degeneration in the CNS in both rodents and primates. Dornan et al (1991) reported that subcutaneous injections of MDMA (40mg/kg) every 12 hours for 4 consecutive days to sexually vigorous male rats produced a transient disruption in the expression of male sexual behavior when compared to saline injected controls. One week after the first injection, MDMA induced a disruption in the expression of male rat copulatory behavior when compared to saline injected controls. In MDMA treated males that did display ejaculatory behavior, however, both the ejaculation latency and the post-ejaculatory interval were dramatically lengthened when compared to controls. This inhibition was not seen, however, at a lower dose of 20mg/kg. One week after the first behavioral test, sexual behavior in MDMA treated rats appeared unaffected despite a marked depletion of 5-HT and 5-HIAA content. Presently, little is known about the effects on the expression of male rat copulatory behavior following administration of other amphetamines which are selective 5-HT neurotoxins. In this study, the effects on male rat copulatory behavior of parachloroamphetamine (PCA), a neurotoxin similar to MDMA, was examined. PCA (10mg/kg and 20mg/kg) or saline (1ml/kg) were administered intraperitoneally to sexually vigorous male rats. These doses of PCA have been shown to produce 5-HT neurotoxicity in the CNS of the rodent. Three days following the injection, a motor activity test was scored and a variety of parameters of male sexual behavior were assessed. One week later the males were tested again. The results of this study revealed that neither neurotoxic doses (10mg/kg and 20mg/kg) of the 5-HT neurotoxin PCA produces the transient disruption of male rat copulatory behavior seen when the similar 5-HT neurotoxin MDMA is administered on a neurotoxic regimen.